EAST Search History

Ref #	Hits	Search Query	DBs	Default Operator	Plurals	Time Stamp
L1	2	(hydrophobin) and (plastic-degrading enzyme)	US-PGPUB; USPAT; USOCR; FPRS; EPO; JPO; DERWENT	ADJ	ON	2007/06/26 09:47
L2	190	(hydrophobin)	US-PGPUB; USPAT; USOCR; FPRS; EPO; JPO; DERWENT	ADJ	ON	2007/06/26 09:42
L3	46	L2 and @py<"2003"	US-PGPUB; USPAT; USOCR; FPRS; EPO; JPO; DERWENT	ADJ	ON	2007/06/26 09:42
L4	9	(plastic-degrading enzyme)	US-PGPUB; USPAT; USOCR; FPRS; EPO; JPO; DERWENT	ADJ	ON .	2007/06/26 09:49
L5	3	"6902887".pn.	US-PGPUB; USPAT; USOCR; FPRS; EPO; JPO; DERWENT	ADJ	ON	2007/06/26 10:02
L6	11050	(abe\$.in.)	USPAT	ADJ	ON	2007/06/26 10:05
L7	0	6 and (degrading plastic)	USPAT	ADJ	ON	2007/06/26 10:06
L8	1483	6 and (plastic)	USPAT ·	ADJ	ON	2007/06/26 10:06
L9	1	8 and (hydrophobin)	USPAT	ADJ	ON	2007/06/26 10:07
L10	410	gomi\$.in.	USPAT	ADJ	ON	2007/06/26 10:07
L11	56	10 and (plastic)	USPAT	ADJ	ON	2007/06/26 10:08
L12	0	11 and (hydrophobin)	USPAT	ADJ	ON	2007/06/26 10:08

6/26/2007 10:14:10 AM Page 1

SCITUS for scientific information only



About Us

Newsroom

Advisory Board

Submit Web Site

Help

Contact Us

	Basic Sea	arch		Advanced Search Search	<u>ch Preferences</u>			
		"hydrophobin"			Search	• •		
	<u>~</u>	Journal sources	✓ Preferred Web sources	Other Web sources	Exact phrase			
Sea	arched for::	:All of the words	:"hydrophobin"					
	Found::	:540 total <u>27</u>	2 journal results 84 pre	ferred web results 1	84 other web result	<u>:s</u>		
	Sort by::	:relevance <u>da</u>	nte					
	Save check	ed results	Email checked results	Export checked resul	ts	Re		
21.		of hydrophobin	n 1 gene transcript in de	eveloping fruiting bod	ies of Lentinula	us foi		
	edodes.	Hiroo / Miva	zaki Yasumasa / Kar	neko. Shinva / Shis	hido. Kazuo .	aeı		
	Nishizawa, Hiroo / Miyazaki, Yasumasa / Kaneko, Shinya / Shishido, Kazuo , Bioscience, biotechnology, and biochemistry, 66 (9), p.1951-1954, Sep 2002							
	Results of in situ RNA-RNA hybridization showed the presence of transcripts of the Lentinula edodes hydrophobin 1 gene, Le.hyd1, everywhere in the mycelial tissues of developing fruiting bodies except for the top parts of the pileus (cap							
	0							
	MEDLINE/	PubMed Citati	on on Pub Med			<u>fur</u>		
	view all 117	results from M	IEDLINE/PubMed			fur gei		
	<u>similar results</u>							
22.	Expression of a fungal hydrophobin in the Saccharomyces cerevisiae cell wall: effect on							
	<u>cell surface properties and immobilization.</u> Nakari-Setälä, Tiina / Azeredo, Joana / Henriques, Mariana / Oliveira, Rosário /							
	Teixeira, José / Linder, Markus / Penttilä, Merja , Applied and environmental							
	microbiology, 68 (7), p.3385-3391, Jul 2002cerevisiae by expression of the HFBI hydrophobin of the filamentous fungus							
	cerevisiae	e by expression	of the HFBI hydropho l the binding affinity of th	bin of the filamentou	is fungus ducing yeast to	sch		
	hvdrophobi	aincrease in c c silicone-based	lin the initial adsorption	on rate of the hydro	phobin yeast was	sec am		
	observed. Furthermore, due							
	MEDLINE/	PubMed Citati	ion on Pub Med			<u>tef</u> tric		
			·			Or		
	similar resu		IEDLINE/PubMed			A		
22			n method with detergen	t/nolymer systems fr	or primary			
<u> </u>	A novel two-step extraction method with detergent/polymer systems for primary recovery of the fusion protein endoglucanase I-hydrophobin I. Collén, Anna / Persson, Josefine / Linder, Markus / Nakari-Setälä, Tiina / Penttilä, Merja / Tjerneld, Folke / Sivars, Ulf , Biochimica et biophysica acta, 1569							
	membrar amphiphilic	fusion protein	2 e, we examine the part endoglucanase I(core)-l from a Trichoderma ree	hydrophobin I (EGI	(core)-HFBI) from			

system...

MEDLINE/PubMed Citation on Publimed

view all 117 results from MEDLINE/PubMed similar results

24. Immunoglobulin E-binding and skin test reactivity to hydrophobin HCh-1 from Cladosporium herbarum, the first allergenic cell wall component of fungi.

Weichel, M / Schmid-Grendelmeier, P / Rhyner, C / Achatz, G / Blaser, K / Crameri, R , Clinical and experimental allergy : journal of the British Society for Allergy and Clinical Immunology, 33 (1), p.72-77, Jan 2003

...classical signature of members of the hydrophobin family. The recombinant protein, termed...the allergenic nature of C. herbarum hydrophobin and indicating a prevalence of sensitization...the range of 8-9%. In contrast, the hydrophobin HYP1 from Aspergillus fumigatus was...

MEDLINE/PubMed Citation on Pub Med

view all 117 results from MEDLINE/PubMed similar results

25. Overproduction, purification, and characterization of the Trichoderma reesei hydrophobin

Askolin, S / Nakari-Setälä, T / Tenkanen, M , Applied microbiology and biotechnology, 57 (1-2), p.124-130, Oct 2001

Many characteristics of fungal hydrophobins, such as an ability to change hydrophobicity...production and purification of a hydrophobin, HFBI of Trichoderma reesei. A high production level of the class II hydrophobin (0.6 g l(-1)) was obtained by constructing...

MEDLINE/PubMed Citation on Publimed

view all 117 results from MEDLINE/PubMed similar results

26. Identification of a hydrophobin gene that is developmentally regulated in the ectomycorrhizal fungus Tricholoma terreum.

Mankel, Angela / Krause, Katrin / Kothe, Erika , Applied and environmental microbiology, 68 (3), p.1408-1413, Mar 2002

...this process. We determined the role of hydrophobins produced by Tricholoma terreum in mycorrhiza formation and hyphal development. A hydrophobin was purified from culture supernatant...heterologous antiserum directed against a hydrophobin found in the aerial mycelium of Schizophyllum...

MEDLINE/PubMed Citation on Pub Med

view all 117 results from MEDLINE/PubMed similar results

27. Cloning and expression analysis of a new hydrophobin cDNA from the ectomycorrhizal basidiomycete Pisolithus.

Duplessis, S / Sorin, C / Voiblet, C / Palin, B / Martin, F / Tagu, D , Current genetics, 39 (5-6), p.335-339, Jul 2001

Hydrophobins are fungal cell wall proteins which play a crucial...and aggregative processes. We have identified a new hydrophobin cDNA (hydPt-3) in the symbiotic mycelium of Pisolithus...previously identified Pisolithus symbiosis-regulated hydrophobins, hydPt-1 and hydPt-2. Also, expression analyses demonstrated...

MEDLINE/PubMed Citation on Pub

view all 117 results from MEDLINE/PubMed similar results

_ 2	28.	A novel two-step extraction method with detergent/polymer systems for primary recovery of the fusion protein endoglucanase Collén, Anna / Persson, Josefine / Linder, Markus / Nakari-Setälä, Tiina / Penttilä, Merja / Tjerneld, Folke / Sivars, Ulf, article, Jan 2002 Extraction systems for hydrophobically tagged proteins have been developed based on phase separation in aqueous solutions of non-ionic detergents and polymers. The systems have earlier only been applied for separation of membrane proteins. Here, we [http://lu-research.lub.lu.se/php/gateway.php?who=lr&me] view all 3 results from Digital Archives similar results
_ 2	9.	The hydrophobin EAS is largely unstructured in solution and functions by forming amyloid-like structures. Mackay, J P / Matthews, J M / Winefield, R D / Mackay, L G / Haverkamp, R G / Templeton, M D , Structure (London, England: 1993), 9 (2), p.83-91, Feb 2001 BACKGROUND: Fungal hydrophobin proteins have the remarkable abilityelusive. We have studied EAS, the hydrophobin from the ascomycete Neurospora crassaunderstand the structural aspects of hydrophobin polymerization. RESULTS: We have
		MEDLINE/PubMed Citation on Pub Med
		view all 117 results from MEDLINE/PubMed similar results
<u> </u>	0.	Cerato-ulmin, a toxin involved in Dutch elm disease, is a fungal hydrophobin. Stringer, M A / Timberlake, W E , The Plant Cell, Jul 2003
		Published journal article available from PubMed Central
		view all 20 results from Pubmed Central similar results
3	1.	Method of binding a compound to a sensor surface using hydrophobin Rogalska, Eva Maria / Tagu, Denis Etienne Marie André / Bilewicz, Renata, EUROPEAN PATENT APPLICATION, Jan 2003 patno: EP1279742
		method comprising the step of adsorbing hydrophobin to said sensor surface. Generally, thereachieve this, it is known to adsorb hydrophobin to the surface of a sensor, and to chemically link a compound to the adsorbed hydrophobin . In particular, Wessels et al. (Advances
•		Full text available at patent office. For more in-depth searching go to view all 69 results from Patent Offices similar results
<u> </u>	2.	Bioscience, Biotechnology and Biochemistry [23K] Jul 2003
		Toluene In Seon KIM,1, Jae Han SHIM,2 and Yong Tack SUH2 p.1945 Note Distribution of Hydrophobin 1 Gene Transcript in Developing Fruiting Bodies of Lentinula edodes Hiroo NISHIZAWA,a Yasumasa [http://www.jsbba.or.jp/e/e_05/bbb6609e.html] similar results
<u> </u>	3.	Hydrophobin gene expression affects hyphal wall composition in Schizophyllum commune.
		van Wetter, M A / Wösten, H A / Sietsma, J H / Wessels, J G, Fungal genetics and biology: FG & B, 31 (2), p.99-104, Nov 2000 Disruption of the SC3 hydrophobin gene of Schizophyllum communeReintroduction of the SC3 gene or other hydrophobins genes expressed behind the SC3relatively high. These data show that hydrophobins not only function at hydrophilic

MEDLINE/PubMed Citation on Pub Med

view all 117 results from MEDLINE/PubMed similar results

34. A gene encoding a hydrophobin, fvh1, is specifically expressed after the induction of fruiting in the edible mushroom Flammulina velutipes.

Ando, A / Harada, A / Miura, K / Tamai, Y , Current genetics, 39 (3), p.190-197, May 2001

...velutipes. Sequence analysis showed that fvh1 encoded for a **hydrophobin**, a small fungal protein usually secreted by filamentous fungi...sequence and a hydropathy pattern characteristics of class I **hydrophobin**. A genomic fvh1 clone was isolated from a F. velutipes genomic...

MEDLINE/PubMed Citation on Pub Med

view all 117 results from MEDLINE/PubMed similar results

35. The hydrophobin HCf-1 of Cladosporium fulvum is required for efficient water-mediated dispersal of conidia.

Whiteford, J R / Spanu, P D, Fungal genetics and biology: FG & B, 32 (3), p.159-168, Apr 2001

Six **hydrophobin** genes (HCf-1 to -6) have thus far been identified in...pathogen Cladosporium fulvum. HCf-1 to -4 are Class I **hydrophobins** and HCf-5 and -6 are Class II **hydrophobins**. In this paper we describe the isolation of deletion...

MEDLINE/PubMed Citation on Pub Med

view all 117 results from MEDLINE/PubMed similar results

36. Spectroscopic Evidence for Amyloid-like Interfacial Self-Assembly of Hydrophobin Sc3
Butko, P. / Buford, J.P. / Goodwin, J.S. / Stroud, P.A. / McCormick, C.L. /
Cannon, G.C. , Biochemical and Biophysical Research Communications, 280 (1), p.212-215, Jan 2001

...Amyloid-like Interfacial Self-Assembly of **Hydrophobin** Sc3 Peter Butko Justin P. Buford 1 1...Congo red REFERENCES 1 Wessels J. G. H. **Hydrophobins**: Proteins that change the nature of...G. H. Interfacial self-assembly of a **hydrophobin** into an amphipathic protein membrane...

Published journal article available from ScienceDirect

view all 134 results from ScienceDirect similar results

37. HCf-6, a novel class II hydrophobin from Cladosporium fulvum

Nielsen, P.S. / Clark, A.J. / Oliver, R.P. / Huber, M. / Spanu, P.D., Microbiological Research, 156 (1), p.59-63, Jan 2001

...Fischer Verlag HCf-6, a novel class II **hydrophobin** from Cladosporium fulvum Peter S. Nielsen...been shown to express a complex family of **hydrophobin** genes including four class I **hydrophobins** and one class II **hydrophobin**. Here we describe...

Published journal article available from ScienceDirect

<u>view all 134 results from ScienceDirect</u> <u>similar results</u>

38. Structural and functional role of the disulfide bridges in the hydrophobin SC3.

de Vocht, M L / Reviakine, I / Wösten, H A / Brisson, A / Wessels, J G /
Robillard, G T , The Journal of biological chemistry, 275 (37), p.28428-28432, Sep 2000

Hydrophobins function in fungal development by self-assembly at hydrophobic-hydrophilic...bridges on the self-assembly, the disulfides of the SC3 **hydrophobin** were

reduced with 1,4-dithiothreitol.	. The free thiolsare not	directly involved in	self-assembly
but keep hydrophobin monon	ners soluble in the funga	I cell or its aqueous	environment

MEDLINE/PubMed Citation on Pub Med

view all 117 results from MEDLINE/PubMed similar results

39. HCf-6, a novel class II hydrophobin from Cladosporium fulvum.

Nielsen, PS / Clark, AJ / Oliver, RP / Huber, M / Spanu, PD, Microbiological research, 156 (1), p.59-63, Jan 2001

...fungal tomato pathogen, has previously been shown to express a complex family of **hydrophobin** genes including four class I **hydrophobins** and one class II **hydrophobin**. Here we describe a gene for HCf-6, a sixth member of the **hydrophobin** family and...

MEDLINE/PubMed Citation on Pub Med

view all 117 results from MEDLINE/PubMed similar results

40. The Hydrophobin HCf-1 of Cladosporium fulvum Is Required for Efficient Water-Mediated Dispersal of Conidia

Whiteford, J.R. / Spanu, P.D., Fungal Genetics and Biology, 32 (3), p.159-168, Apr 2001

...91263-0 Academic Press Regular Article The **Hydrophobin** HCf-1 of Cladosporium fulvum Is Required...Medicine, London, United Kingdom Six **hydrophobin** genes (HCf-1 to -6) have thus far...Cladosporium fulvum. HCf-1 to -4 are Class I **hydrophobins** and HCf-5 and -6 are Class II **hydrophobins**...

Published journal article available from ScienceDirect

view all 134 results from ScienceDirect similar results

Sponsored links

Hydrophobin

Relax. Take a deep breath. We have the answers you seek.

www.righthealth.com

:::fast

Results Pages: [<< Prev] 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 [Next >>] back to to

"hydrophobin"		Search	
✓ lournal sources	✓ Preferred Web sources	Other Web sources	Exact phrase

<u>Downloads</u> | <u>Library Partners</u> | <u>Subscribe to News Updates</u> | <u>User Feedback</u> <u>Advertising</u> | <u>Tell A Friend</u> | <u>Terms Of Service</u> | <u>Privacy Policy</u> | <u>Legal</u>

Powered by FAST © Elsevier 2007



Video News Maps "hydrophobin"

- 2003

Search

Scholar All articles - Recent articles Results 1 - 10 of about 642 for "hydrophobin". (0.08 seconds)

All Results

Interfacial Self-Assembly of a Fungal Hydrophobin into a Hydrophobic

J Wessels

Rodlet Layer - all 5 versions »

HAB Wosten, OMH de Vries, JGH Wessels - The Plant Cell Online, 1993 - Am Soc Plant Biol

O de Vries ... RESEARCH ARTICLES. Interfacial Self-Assembly of a Fungal Hydrophobin into a

H Wösten

Hydrophobic Rodlet Layer. HAB. Wosten, OMH. de Vries and JGH. ...

H Wosten

Cited by 87 - Related Articles - Web Search - BL Direct

F Schuren

Interfacial self-assembly of a hydrophobin into an amphipathic protein membrane mediates fungal ... - all 3 versions »

HAB Wösten, FHJ Schuren, JGH Wessels - EMBO J, 1994 - pubmedcentral.nih.gov ... Copyright notice. Interfacial self-assembly of a hydrophobin into an amphipathic protein membrane mediates fungal attachment to hydrophobic surfaces. ...

Cited by 94 - Related Articles - Web Search - BL Direct

An abundant hydrophobin (ABH1) forms hydrophobic rodlet layers in Agaricus bisporus fruiting bodies - all 5 versions »

LG Lugones, JS Bosscher, K Scholtmeyer, OM de ... - Microbiology, 1996 - Soc General Microbiol

... Microbiology. ARTICLES. An abundant hydrophobin (ABH1) forms hydrophobic rodlet layers in Agaricus bisporus fruiting bodies. LG Lugones ...

Cited by 52 - Related Articles - Web Search - BL Direct

MPG1 Encodes a Fungal Hydrophobin Involved in Surface Interactions during Infection-Related ... - all 6 versions »

NJ Talbot, MJ Kershaw, GE Wakley, OMH de Vries, ... - The Plant Cell Online, 1996 - Am Soc Plant Biol

... MPGI Encodes a Funga1 Hydrophobin Involved in Surface Interactions during Infection-Related Development of ... MPG1 Directs Formation of a Class I Hydrophobin ... Cited by 77 - Related Articles - Web Search - BL Direct

... relationships in Aspergillus section Fumigati inferred from partial betatubulin and hydrophobin DNA ... - all 4 versions »

DM Geiser, JC Frisvad, JW Taylor - Mycologia, 1998 - apt.allenpress.com 90, No. 5, pp. 831-845. Evolutionary relationships in Aspergillus section Fumigati inferred from partial β-tubulin and hydrophobin DNA sequences. ...

Cited by 58 - Related Articles - Web Search - BL Direct

MPG1, a gene encoding a fungal hydrophobin of Magnaporthe grisea, is involved in surface recognition ... - all 2 versions »

JL Beckerman, DJ Ebbole - Mol Plant Microbe Interact, 1996 - ncbi.nlm.nih.gov MPG1, a gene encoding a fungal hydrophobin of Magnaporthe grisea, is involved in surface recognition. Beckerman JL, Ebbole DJ. Department ...

Cited by 53 - Related Articles - Web Search - BL Direct

Hydrophobin Genes Involved in Formation of Aerial Hyphae and Fruit Bodies in Schizophyllum - all 5 versions »

JGH Wessels, OMH de Vries, SA Asgeirsdottir, FHJ ... - The Plant Cell Online, 1991 - Am Soc Plant Biol

... Physiologists **Hydrophobin** Genes Involved in Formation of Aerial ... hyphae. The Sc3 gene encodes a **hydrophobin** present in walls of aerial hyphae. ...

Cited by 61 - Related Articles - Web Search

Insoluble hydrophobin complexes in the walls of Schizophyllum commune and other filamentous fungi - all 2 versions »

OMH Vries, MP Fekkes, HAB Wösten, JGH Wessels - Archives of Microbiology, 1993 - Springer

... Insoluble hydrophobin complexes in the walls of ... 1), previously shown to be the products

of the Sc3 and Sc4 hydrophobin genes, respectively (Wessels et al. ...

Cited by 56 - Related Articles - Web Search - BL Direct

... -controlled gene, ccg-2, is allelic to eas and encodes a fungal hydrophobin required for formation ... - all 5 versions »

D Bell-Pedersen, JC Dunlap, JJ Loros - Genes and Development, 1992 - Cold Spring Harbor Lab

... The Neurospora circadian clock-controlled gene, ccg-2, is allelic to eas and encodes a fungal **hydrophobin** required for formation of the conidial rodlet layer. ... Cited by 57 - Related Articles - Web Search - BL Direct

Targeted mutation of the SC3 hydrophobin gene of Schizophyllum commune affects formation of aerial ... - all 2 versions »

MA van Wetter, FHJ Schuren, TA Schuurs, JGH ... - FEMS Microbiol. Lett, 1996 - Blackwell Synergy

... Targeted mutation of the SC3 **hydrophobin** gene of ... The SC3 **hydrophobin** gene of Schizophyllum commune was disrupted by homologous integration of an SC3 genomic ... Cited by 47 - Related Articles - Web Search

Goo	000	000	oog	1	e	

Result Page: 1 2 3 4 5 6 7 8 9 10 Next

		 _	
'hydr	ophobin"		Search

Google Home - About Google - About Google Scholar

©2007 Google

1996 - Am Soc Microbiol



<u>Video News Maps more »</u>

"Cutinase" and "Aspergillus"

- 2003

Search

The "AND" operator is unnecessary -- we include all search terms by default. [details]

Scholar All articles - Recent articles Results 31 - 40 of about 490 for "Cutinase" and "Aspergillus"

All Results

C Verrips

T Cleveland

R Gouka

W Musters J Sweigard

Cloning of Cutinase Transcription Factor 1, a Transactivating Protein Containing Cys6Zn2 Binuclear ... - all 3 versions »

... pectinase (1 g/liter) and Aspergillus niger cellulase (5 g/liter; Sigma Chemical

Co ... Because the cutinase-negative mutant does not grow on cutin, cultures to ...

D Li, PE Kolattukudy - Journal of Biological Chemistry, 1997 - ASBMB

... cDNA clone encoding a polypeptide designated cutinase transcription factor 1 ... carlsbergensis, Kluyveromyces lactis, Neurospora crassa, Aspergillus nidulans, and ...

Fusarium polycaprolactone depolymerase is cutinase - all 4 versions »

CA Murphy, JA Cameron, SJ Huang, RT Vinopal - Applied and Environmental Microbiology,

Cited by 22 - Related Articles - Web Search - BL Direct

Cited by 39 - Related Articles - Web Search - BL Direct

Surface-induced changes in the structure and activity of enzymes physically immobilized at solid/ ... - all 9 versions »

W Norde, T Zoungrana - Biotechnology and Applied Biochemistry, 1998 bab.portlandpress.com

... Cutinase from Aspergillus oryzae was a gift from Unilever (Vlaardingen, The Netherlands). Cutinase is also able to hydrolyse soluble esters. Page 3. ... Cited by 36 - Related Articles - View as HTML - Web Search - BL Direct

Inactivation of polycaprolactone depolymerase (cutinase) in Fusarium cultures by an extracellular ... - all 3 versions »

CA Murphy, JA Cameron, SJ Huang, RT Vinopal - Journal of Industrial Microbiology and Biotechnology, 1999 - Springer

... Because the opti- mum pH for Fusarium cutinase activity is between 9 and ... pectin lyase B and secreted heterologous cloned proteins in Aspergillus niger cultures ... Cited by 1 - Related Articles - Web Search - BL Direct

Effect of disruption of a cutinase gene (cutA) on virulence and tissue specificity of Fusarium ... - all 6 versions »

RN Crowhurst, SJ Binnie, JK Bowen, BT Hawthorne, ... - Mol Plant Microbe Interact, 1997 apsnet.org

... Genes encoding cutinase have been cloned from several fungi including Alternaria brassicicola (Yao and Köller 1994), Aspergillus oryzae (Ohnishi et al. ...

Cited by 13 - Related Articles - View as HTML - Web Search - BL Direct

Induction of a Biopolyester Hydrolase (Cutinase) by Low Levels of Cutin Monomers in Fusarium solani ... - all 3 versions »

TS Lin, PE Kolattukudy - Journal of Bacteriology, 1978 - Am Soc Microbiol ... in the medium, thus showing a dual control of cutinase production. ... p-nitrophenyl butyrate (PNB), o- dianisidine, glucose oxidase (from Aspergillus niger), and ... Cited by 26 - Related Articles - Web Search

Studies on ferulic acid esterase activity in fungal lipases and cutinases - all 3 versions »

A Andersen, A Svendsen, J Vind, SF Lassen, C Hjort ... - Colloids and Surfaces B: Biointerfaces, 2002 - aapspharmaceutica.com

... In addition to lipase activity, three cutinases showed ferulic acid esterase activity, Aspergillus oryzae cutinase (5 U/mg), Fusarium solani pisi cutinase (13 U ...

Cited by 7 - Related Articles - Cached - Web Search

Regulation of Constitutively Expressed and Induced Cutinase Genes by Different Zinc Finger ... - all 5 versions »

D Li, T Sirakova, L Rogers, WF Ettinger, PE ... - Journal of Biological Chemistry, 2002 - ASBMB

... Binding to the palindromic DNA element in the **cutinase** promoter by the expressed ... NIRA (39), UAY (40), QUTA (41), and AMDR (42) of **Aspergillus** nidulans; and ... Cited by 11 - Related Articles - Web Search - BL Direct

... the cDNA and gene for an elastinolytic aspartic proteinase from.

Aspergillus fumigatus and evidence ... - all 5 versions »

JD Lee, PE Kolattukudy - Infection and Immunity, 1995 - Am Soc Microbiol ... and deletion of the gene encoding aspergillopepsin A from **Aspergillus** awamori. ... Identifi-

cation of regulatory elements in the **cutinase** promoter from Fusarium ... Cited by 26 - Related Articles - Web Search - BL Direct


Can hTNF-a be successfully produced and secreted in filamentous fungus

Aspergillus niger?

N Kraševec, CA van den Hondel, R Komel - Pflügers Archiv European Journal of Physiology, 2000 - Springer

... in the GRAS filamentous fungus A. niger, because the **Aspergillus** expression-secretion ...

dserent efficiencies, as for example in the case of **cutinase** mutants [III ... Web Search



Result Page: <u>Previous 1 2 3 4 5 6 7 8 9 10111213</u> Next

"Cutinase" and "Aspergillus" Search

Google Home - About Google - About Google Scholar

©2007 Google